

LEVEL STATISTICS IN THE QUANTUM HALL REGIME

M. BATSCCH^{1,2} and L. SCHWEITZER¹

¹*Physikalisch-Technische Bundesanstalt, Bundesallee 100,
D-38116 Braunschweig, Germany*

²*I. Institut für Theoretische Physik, Jungiusstraße 9,
D-20355 Hamburg, Germany*

The statistical properties of energy eigenvalues in the critical regime of the lowest Landau band are investigated. The nearest neighbour level spacing distribution $P(s)$ and the Mehta quantity I_0 , from which the scaling exponent ν of the correlation length can be derived, are calculated. While $P(s)$ disagrees with recent analytical predictions, ν is found to be in agreement with theoretical and experimental results.

1 Introduction

In recent years the investigation of energy level statistics in systems showing a localisation-delocalisation transition has made remarkable progress. For the metallic side of the transition the spectral fluctuations only depend on the symmetry of the system (orthogonal, unitary or symplectic) and are well described by Random Matrix Theory (RMT)¹. On the insulating side, the energy levels are uncorrelated and exhibit Poissonian behaviour. At the transition point, new critical ensembles were found for the 3d orthogonal^{2,3,4} the 2d symplectic^{5,6}, and the 3d unitary case⁷. In the quantum Hall system no complete metal-insulator transition is observed, and the situation is unclear^{8,9,10,11}.

This is due to the fact that the localisation length of the electronic wavefunctions diverges at the centre of the disorder broadened Landau bands. Hence, in the thermodynamical limit there is no range of extended states but only one energy where the states are multifractal^{8,12,13}. Therefore, the question is whether there exists a scale-invariant level statistics in *finite systems* in the absence of true extended states.

As a result of analytical theories^{14,15} the asymptotic behaviour of the nearest neighbour level spacing distribution $P(s)$, where the energy difference s is measured in units of the mean level spacing Δ , was proposed to obey $P(s) \propto \exp(-cs^{2-\gamma})$ with $\gamma = 1 - \frac{1}{\nu d}$, where ν is the critical exponent and d the Euclidean dimension. Fits to the numerical data in 3d orthogonal⁴, 3d unitary⁷ and 2d symplectic^{5,6} systems did not result in a sufficient agreement. Investigations that seem to show a reasonable fit^{11,16} require, however, values for ν which are different from those commonly accepted. Previously, a simple exponential decay of $P(s)$ was proposed², $P(s) \propto \exp(-\kappa s)$, where $\kappa > 1$ is a




Figure 1: Energy and size dependence of I_0 at the critical energy in the lowest Landau band

numerical constant. From the calculations in 3d orthogonal⁴ and 3d unitary systems⁷, $\kappa \approx 1.9$, and for the 2d symplectic, $\kappa \approx 4.0$ was obtained⁵. In our opinion, κ should reflect the fractal structure of the critical wavefunctions. Therefore, it is very interesting to study the asymptotics of $P(s)$ in the 2d QHE situation where no complete localisation-delocalisation transition occurs.

2 Model and method

We investigate a 2d Anderson model with the magnetic field included via a phase-factor in the transfer matrix elements V (see Ref⁸ for details). The diagonal disorder potentials are taken at random with constant probability from $[-W/2, W/2]$ and $W/V = 2.5$. The strength of the magnetic field equals $1/8$ flux quanta per unit cell. We diagonalise the corresponding Hamiltonian matrix for finite squares of size $L^2 = 32^2, 64^2, 128^2$, and 160^2 using a Lanczos procedure. To obtain a reliable statistic, up to 3^6 realizations were calculated.

3 Results and discussion

To find the energy range of the eigenstates with localisation length larger than the system size and to extract the critical exponent ν , we employ the quantity




Figure 2: The critical QHE- $P(s)$ ($\bullet$) in comparison with the critical 3d unitary ($\star$), the Poisson and the GUE result. The inset shows a $\exp(-\kappa s)$ decay with $\kappa \approx 4.1$

I_0 known from RMT¹. The calculation does neither require a fitting procedure as the Brody level repulsion parameter⁹, nor some cutoff value, as for example the integrated Δ_3 statistics¹⁷. I_0 is obtained by integrating over $P(s)$

$$I_0 = \int_0^\infty \int_s^\infty \int_{s'}^\infty P(s'') ds'' ds' ds. \quad (1)$$

The value of I_0 for the Gaussian unitary ensemble is $I_0 = 0.590$ and for uncorrelated (localised) levels $I_0 = 1$. Multifractal states are expected to have a value larger than 0.59. From our calculations we find $I_0 = 0.605 \pm 0.002$ near the critical energy independent of the considered system sizes. Fig. 1 displays the energy dependence of I_0 in the energy range about the critical point. We have carefully checked that I_0 does not depend on the width of the energy interval from which the eigenvalues are taken. For all system sizes one observes a minimum $I_0 \approx 0.605$ at the critical energy $E_c/V = -3.33$. States away from E_c become more localised when the system size is enlarged. From the scaling behaviour of $I_0(E, L)$ we extract the critical exponent $\nu = 2.4$ in good agreement with both theoretical¹³ and experimental¹⁸ results.

The nearest neighbour level spacing distribution $P(s)$ is shown in Fig. 2. It clearly deviates from the Wigner surmise although it shows a quadratic small- s behaviour as expected for a unitary symmetry. Our result for the lattice model

resembles that obtained for a continuum model^{9,11}. Due to the large number of realizations we are able to extract the large- s behaviour displayed in the inset of Fig. 2. As for the other critical ensembles, it is not possible to fit the tail of $P(s)$ by the proposal¹⁵ $P(s) \sim \exp(-cs^{2-\gamma})$ with $\nu = 2.4$, but it can be fitted by a simple exponential decay, $P(s) \propto \exp(-\kappa s)$, with $\kappa \approx 4.1$. This value is similar to the one found for a 2d system with symplectic symmetry⁵. In conclusion, there exists a critical level statistics in 2d QHE systems despite the absence of a complete metal-insulator transition.

References

1. M. L. Metha, *Random Matrices*, 2nd ed. (Academic Press, San Diego, 1991).
2. B. L. Altshuler, I. Kh. Zharekeshev, S. A. Kotochigova, and B. Shklovskii, Sov. Phys. JETP **67**, 625 (1988).
3. B. I. Shklovskii *et al.*, Phys. Rev. B **47**, 11487 (1993).
4. I. Kh. Zharekeshev and B. Kramer, Jap. J. Appl. Phys. **34**, 4361 (1995).
5. L. Schweitzer and I. Kh. Zharekeshev, J. Phys.: Condens. Matter **7**, L377 (1995).
6. S. N. Evangelou, Phys. Rev. Lett. **75**, 2550 (1995).
7. M. Batsch, L. Schweitzer, I. Kh. Zharekeshev, and B. Kramer, Phys. Rev. Lett. (1996).
8. B. Huckestein and L. Schweitzer, in *High Magnetic Fields in Semiconductor Physics III: Würzburg 1990*, edited by G. Landwehr (Springer Series in Solid-State Sciences 101, Springer, Berlin, 1992), p. 84.
9. Y. Ono, in *High Magnetic Fields in the Physics of Semiconductors*, edited by D. Heiman (World Scientific Publ., Singapore, 1995), p. 260.
10. M. Feingold, Y. Avishai, and R. Berkovits, Phys. Rev. B **52**, 8400 (1995).
11. T. Ohtsuki and Y. Ono, J. Phys. Soc. Jpn. **64**, 4088 (1995).
12. W. Pook and M. Janßen, Z. Phys. B **82**, 295 (1991).
13. B. Huckestein, B. Kramer, and L. Schweitzer, Surf. Science **263**, 125 (1992).
14. V. E. Kravtsov, I. V. Lerner, B. L. Altshuler, and A. G. Aronov, Phys. Rev. Lett. **72**, 888 (1994).
15. A. G. Aronov, V. E. Kravtsov, and I. V. Lerner, JETP Lett. **59**, 39 (1994).
16. S. N. Evangelou, Phys. Rev. B **49**, 16805 (1994).
17. B. Huckestein and L. Schweitzer, Phys. Rev. Lett. **72**, 713 (1994).
18. S. Koch, R. J. Haug, K. v. Klitzing, and K. Ploog, Phys. Rev. Lett. **67**, 883 (1991).